

From [rxpgnews.com](http://rxpgnews.com)

## NEUROSCIENCES

### Artificial synapses created between nanoelectronic devices and mammalian neurons

By Harvard University

Aug 25, 2006, 19:26

#### [Test1](#)

Visit the Red Planet in style. Low-gravity fun for everyone!

[www.fillz.com/](http://www.fillz.com/)

#### [Intraoperative Monitoring](#)

Intraoperative monitoring services for hospitals and surgeons

[www.iomservices.com](http://www.iomservices.com)

[Ads by Goooooogle](#)

Opening a whole new interface between nanotechnology and neuroscience, scientists at Harvard University have used slender silicon nanowires to detect, stimulate, and inhibit nerve signals along the axons and dendrites of live mammalian neurons.

"We describe the first artificial synapses between nanoelectronic devices and individual mammalian neurons, and also the first linking of a solid-state device -- a nanowire transistor -- to the neuronal projections that interconnect and carry information in the brain," says Lieber, the Mark Hyman, Jr., Professor of Chemistry in Harvard's Faculty of Arts and Sciences and Division of Engineering and Applied Sciences. "These extremely local devices can detect, stimulate, and inhibit propagation of neuronal signals with a spa-tial resolution unmatched by existing techniques."

Electrophysiological measurements of brain activity play an important role in understanding signal propagation through individual neurons and neuronal networks, but existing technologies are relatively crude: Micropipette electrodes poked into cells are invasive and harmful, and microfabricated electrode arrays are too bulky to detect activity at the level of individual axons and dendrites, the neuronal projections responsible for electrical signal propagation and inter-neuron communication.

By contrast, the tiny nanowire transistors developed by Lieber and colleagues gently touch a neuronal projection to form a hybrid synapse, making them noninvasive, and are thousands of times smaller than the electronics now used to measure brain activity.

Lieber's group has previously shown that nanowires can detect, with great precision, molecular markers indicating the presence of cancer in the body, as well as single viruses. Their latest work takes advantage of the size similarities between ultra-fine silicon nanowires and the axons and dendrites projecting from nerve cells: Nanowires, like neuronal offshoots, are just tens of nanometers in width, making the thin filaments a good match for intercepting nerve signals.

Because the nanowires are so slight -- their contact with a neuron is no more than 20 millionths of a meter in length -- Lieber and colleagues were able to measure and manipulate electrical conductance at as many as 50 locations along a single axon.

The current work involves measurement of signals only within single mammalian neurons; the researchers are now working toward monitoring signaling among larger networks of nerve cells. Lieber says the devices could also eventually be configured to measure or detect neurotransmitters, the chemicals that leap synapses to carry electrical impulses from one neuron to another.

"This work could have a revolutionary impact on science and technology," Lieber says. "It provides a powerful new approach for neuroscience to

study and manipulate signal propagation in neuronal networks at a level unmatched by other techniques; it provides a new paradigm for building sophisticated interfaces between the brain and external neural prosthetics; it represents a new, powerful, and flexible approach for real-time cellular assays useful for drug discovery and other applications; and it opens the possibility for hybrid circuits that couple the strengths of digital nanoelectronic and biological computing components."

All rights reserved by [www.rxpnews.com](http://www.rxpnews.com)