



Functional Group Imaging by Chemical Force Microscopy

C. Daniel Frisbie, Lawrence F. Rozsnyai, Aleksandr Noy, Mark S. Wrighton, Charles M. Lieber

Science, New Series, Volume 265, Issue 5181, Genome Issue (Sep. 30, 1994), 2071-2074.

Stable URL:

<http://links.jstor.org/sici?sici=0036-8075%2819940930%293%3A265%3C2071%3AFGIBCF%3E2.0.CO%3B2>

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.


Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Science is published by The American Association for the Advancement of Science. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at [http://www.jstor.org/journals/aaas](http://www.jstor.org/journals/aaas.html)

Functional Group Imaging by Chemical Force Microscopy

C. Daniel Frisbie, Lawrence F. Rozsnyai, Aleksandr Noy,
Mark S. Wrighton, Charles M. Lieber*

Mapping the spatial arrangement of chemical functional groups and their interactions is of significant importance to problems ranging from lubrication and adhesion to recognition in biological systems. A force microscope has been used to measure the adhesive and friction forces between molecularly modified probe tips and organic monolayers terminating in a lithographically defined pattern of distinct functional groups. The adhesive interactions between simple CH_3/CH_3 , CH_3/COOH , and COOH/COOH functional groups correlate directly with friction images of sample



- between two CH₃ groups, which assumes the interaction can be treated with a Lennard-Jones potential (27).
24. J. H. Hoh, J. P. Cleveland, C. B. Prater, J.-P. Revel, P. K. Hansma, *J. Am. Chem. Soc.* **114**, 4917 (1992).
25. G. Liu and M. B. Salmeron, *Langmuir* **10**, 367 (1994).
26. It has been suggested on the basis of SFA studies of different hydrocarbon-terminated layers that it is the hysteresis in adhesion that will correlate with friction (3). We believe that in comparisons of chemically distinct functional groups, which exhibit large differences in adhesive interaction, friction differences are directly related to the variations in the adhesive forces.
27. A. Noy, C. D. Frisbie, C. M. Lieber, unpublished results.

-